

## Specialty Infusion Apothecary & Wellness Spa INFUSION ORDERS- REMICADE (INFLIXIMAB)

PATIENT INFORMATION				
Name: DOB:				
Allergies: Date of Referral:				
REFERRAL STATUS				
☐ New Referral ☐ Dose or Frequency Change ☐ Order Renewal				
DIAGNOSIS AND ICD 10 CODE				
☐ Moderate to Severe Ulcerative Colitis ICD 10 Code: K51.90				
☐ Moderate to Severe Crohn's Disease ICD 10 Code: K50				
☐ Rheumatoid Arthritis ICD 10 Code: M			6.9	
☐ Ankylosing Spondylitis ICD 10 Code: M45.				
□ Psoriatic Arthritis ICD 10 Code: L40.52				
☐ Plaque Psoriasis ICD 10 Code: L40.0				
☐ Other: ICD10 Code:				
Guiet.				
REQUIRED DOCUMENTATION				
☐ This signed order form by the provider			☐ Clinical/Progress notes	
☐ Patient demographics AND insurance information			□ Labs	and Tests supporting primary diagnosis
☐ Hepatitis B Test Results: HBsAg, Total HepB Core Antibody			☐ TB Test Results	
List Tried & Failed Therapies, including duration of treatment:				
1)				
2)				
3)				
MEDICATION ORDERS				
Initial Dosing	☐ Remicade 5mg/kg IV at week 0, 2, 6, then every 8 weeks thereafter			
Maintenance Dosing	☐ Remicade 5mg/kg IV every 8 weeks			
Alternative Dosing	□ Remicade	IV every	weeks	
atient Weight= kg				
Refills:				
PREMEDICATIONS				
☐ Acetaminophen 650mg PO prior to Remicade infusion				
☐ Diphenhydramine 25mg PO prior to Remicade infusion				
☐ Methylprednisolone 40mg Slow IV Push PRN infusion reaction				
□ Other:				
Please note: if an infusion reaction occurs, the on-call physician will order appropriate rescue medications as deemed medically necessary.				
This may also include pausing, reducing the rate of infusion or discontinuing the medication.				
PRESCRIBER INFORMATION  Dresseriber Names				
Prescriber Name:				Office
Office Phone:	Office Fax:			Office Email:
	Office Pax.			
Prescriber Signature:				Date:

Contact us with questions at: info@siainfusions.com or call (561) 955-7079